


UNIT-III Complex Analysis

Functions of a Complex Variable :-

Complex number $z = x + iy$; $i = \sqrt{-1}$

If (r, θ) be the polar coordinates of P -



$$z = r \cos \theta + i r \sin \theta$$

$$= r [\cos \theta + i \sin \theta]$$

$$\boxed{z = r e^{i\theta}} \Rightarrow \text{represents complex number in Polar form}$$

If $z = x + iy$ is a complex number then $\bar{z} = x - iy$ is a complex conjugate.

NOTE:- $|z - a| = r$ represents a circle with centre at a & radius r .

Function of a complex variable -

If x & y are real variables then $z = x + iy$ is called a complex variable. If corresponding to each value of a complex variable z in a given region R there correspond one or more values of another complex variable $w = [u + iv]$

{ If z is a comp. variable $x + iy \Rightarrow w$ is also comp. i.e. $u + iv$.
 w is called function of complex variable z & is denoted as $w = f(z) = u + iv$.

Derivative of $f(z)$:-

Let $w = f(z)$ be a single valued functⁿ of variable $z = x + iy$ then derivative of $w = f(z)$ is

$$\frac{dw}{dz} = f'(z) = \lim_{\delta z \rightarrow 0} \frac{f(z + \delta z) - f(z)}{\delta z} \text{ provided the}$$

limit exists and has same value for all the diff. ways in which δz approaches to zero.

Analytic Functions -

A function $f(z)$ which is a single valued and possesses a unique derivative at all points of region R then $f(z)$ is called an analytic function in that region.

Remark: i) An analytic functⁿ is also called regular functⁿ or holomorphic functⁿ.

ii) Functⁿ which is analytic everywhere in the complex plane is known as an entire function.

iii) A point at which an analytic functⁿ fails to be analytic or fails to possess a unique derivative is called singular point of function.

Eg: $f(z) = \frac{z}{z-a}$; {functⁿ fails to be analytic at $z=a \therefore z=a$ is a singular pt}

Cauchy-Riemann [C-R] equations in Cartesian form -

(6 Marks) Theorem:

"Necessary conditⁿ that single valued functⁿ $f(z)$.

$f(z) = u(x,y) + i v(x,y)$ be analytic at any pt. $z = x + iy$ is that there exists four continuous first order partial derivatives $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$ which are continuous &

satisfy equations $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ and $\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$, these are

$$(u_x = v_y) \quad (v_x = -u_y)$$

called as CR equations."

⇒ Proof:

Let $f(z)$ be analytic at a point $z \therefore$ by definition of analytic functⁿ

$$f'(z) = \lim_{\Delta z \rightarrow 0} \frac{f(z + \Delta z) - f(z)}{\Delta z} \text{ exists \& is unique.}$$

In cartesian form :-

$$f(z) = u(x,y) + i v(x,y)$$

let $\Delta x, \Delta y, \Delta z$ be the increments in x, y, z respectively

$$\begin{aligned}
 f'(z) &= \lim_{\delta z \rightarrow 0} \frac{[u(x+\delta x, y+\delta y) + i v(x+\delta x, y+\delta y)] - [u(x, y) + i v(x, y)]}{\delta z} \\
 &= \lim_{\delta z \rightarrow 0} \frac{[u(x+\delta x, y+\delta y) - u(x, y)] + i [v(x+\delta x, y+\delta y) - v(x, y)]}{\delta z} \\
 &= \lim_{\delta z \rightarrow 0} \left\{ \frac{u(x+\delta x, y+\delta y) - u(x, y)}{\delta z} + i \frac{[v(x+\delta x, y+\delta y) - v(x, y)]}{\delta z} \right\}
 \end{aligned}$$

since $z = x + iy \therefore \delta z = \delta x + i\delta y$

Since δz approaches 0 in two ways

Case (i): Along real axis {ie along δx }

$$\text{ie } y=0 \Rightarrow \delta y=0$$

$$\Rightarrow \delta z = \delta x$$

Equation (1) becomes-

$$\begin{aligned}
 f'(z) &= \lim_{\delta x \rightarrow 0} \frac{u(x+\delta x, y) - u(x, y)}{\delta x} \\
 &\quad + i \lim_{\delta x \rightarrow 0} \frac{v(x+\delta x, y) - v(x, y)}{\delta x}
 \end{aligned}$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \quad \text{--- (2)}$$

Case (ii): Along imaginary axis:

$$\text{ie } x=0 \Rightarrow \delta x=0$$

$$\Rightarrow \delta z = i\delta y$$

Equation (1) becomes-

$$f'(z) = \lim_{i\delta y \rightarrow 0} \frac{u(x, y+\delta y) - u(x, y)}{i\delta y} + i \lim_{\delta y \rightarrow 0} \frac{v(x, y+\delta y) - v(x, y)}{\delta y}$$

$$f'(z) = \frac{1}{i} \frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \quad \text{--- (3)}$$

$$f'(z) = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \quad \text{--- (3)}$$

Existence of $f'(z)$ thus guarantees the existence of the ^{four} first order partial derivatives which are continuous in eqⁿ (2) & (3)

equating (2) & (3) :-

$$\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y}$$

Equating real part:-

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

Equating imaginary part:-

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

REMARK: The real & imaginary parts of an analytic functⁿ are called conjugate funct^{ns}.

∴ if $f(z) = u(x, y) + iv(x, y)$ is an analytic functⁿ then $u(x, y)$ & $v(x, y)$ are conjugate functions and relation b/w two conjugate functions is given by CR equations.

Couchy - Riemann [C-R] equations in polar form:-

"If a functⁿ $f(z)$ is analytic at a point z then there exists 4 continuous first order partial derivatives

$\frac{\partial u}{\partial r}, \frac{\partial u}{\partial \theta}, \frac{\partial v}{\partial r}, \frac{\partial v}{\partial \theta}$ & satisfy equation called C-R equat^{ns}

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta} \quad \text{and} \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$$

Note:-

$$f'(z) = f'(re^{i\theta})$$

$$= e^{-i\theta} \left[\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right]$$

⇒ Proof:-

let (r, θ) be polar coordinates of pt z whose cartesian coordinates are (x, y) then

$$x = r \cos \theta \quad y = r \sin \theta$$

$$z = x + iy = r \cos \theta + i r \sin \theta$$

$$z = re^{i\theta}$$

$$u + iv = f(z) = f(re^{i\theta}) \quad \text{--- (1)}$$

diff. (1) partially wrt r keeping θ constant

$$\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} = f'(re^{i\theta}) e^{i\theta} \quad \text{--- (2)}$$

diff. (1) partially wrt θ keeping r constant

$$\frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} = r i e^{i\theta} f'(re^{i\theta}) \quad \text{--- (3)}$$

From ② & ③

$$\frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} = ir \left[\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right]$$

equating imaginary parts

$$i \frac{\partial v}{\partial \theta} = ir \frac{\partial u}{\partial r}$$

$$\boxed{\frac{\partial v}{\partial \theta} = r \frac{\partial u}{\partial r}}$$

equating real parts -

$$\boxed{\frac{\partial u}{\partial \theta} = -r \frac{\partial v}{\partial r}}$$

$$\Rightarrow \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta} \quad \& \quad \frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta} \quad \text{are CR equations.}$$

LP3) Prove that given funct^{ns} are analytic & find their derivative.

1) $\sin iz + e^z$

$$f(z) = \sin iz + e^z$$

$$= e^{x+iy}$$

$$= e^x \cdot e^{iy}$$

$$= e^x [\cos y + i \sin y]$$

$$f(z) = e^x \cos y + i e^x \sin y = u + iv$$

$$\Rightarrow u = e^x \cos y \quad v = e^x \sin y$$

$$\frac{\partial u}{\partial x} = e^x \cos y \quad \frac{\partial v}{\partial x} = e^x \sin y$$

$$\frac{\partial u}{\partial y} = -\sin y e^x \quad \frac{\partial v}{\partial y} = \cos y e^x$$

$$\text{since } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \quad \text{CR eq^{ns} are satisfied}$$

$\therefore f(z) = e^z$ is a analytic / holomorphic / regular functⁿ.

$$\begin{aligned}
 f'(z) &= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\
 &= e^x \cos y + i e^x \sin y \\
 &= e^x (\cos y + i \sin y) \\
 &= e^x \cdot e^{iy} \\
 &= e^{x+iy}
 \end{aligned}$$

$$\boxed{f'(z) = e^z}$$

vi) $z + e^z = (x+iy)(e^x \cdot e^{iy})$
 (?) $= (x+iy)(e^x \cdot (\cos y + i \sin y)) = (x+iy)[e^x \cos y + i e^x \sin y]$
 $= x e^x \cos y + i x e^x \sin y + i y e^x \cos y - y e^x \sin y$

$$\begin{aligned}
 f(z) &= [x e^x \cos y - y e^x \sin y] + i [x e^x \sin y + y e^x \cos y] \\
 &= u + i v
 \end{aligned}$$

$$\Rightarrow u = x e^x \cos y - y e^x \sin y \quad \& \quad v = x e^x \sin y + y e^x \cos y$$

$$\frac{\partial u}{\partial x} = \cos y [x e^x + e^x] - e^x y \sin y = e^x [x \cos y + \cos y - y \sin y]$$

$$\frac{\partial u}{\partial y} = -x e^x \sin y - e^x [y \cos y + \sin y]$$

$$\frac{\partial v}{\partial x} = \sin y [x e^x + e^x] + y e^x \cos y$$

$$\frac{\partial v}{\partial y} = x e^x \cos y + e^x [y \sin y + \cos y] = e^x [x \cos y + \cos y - y \sin y]$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y} \quad \text{; CR equations are satisfied.}$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\begin{aligned}
 &= e^x [x \cos y + \cos y - y \sin y] + i [\sin y (x e^x + e^x) + y e^x \cos y] \\
 &= e^x [x \cos y + \cos y - y \sin y + i x \sin y + i \sin y + i y \cos y]
 \end{aligned}$$

$$= e^x [\cos y + x \cos y - y \sin y + i (x \sin y + \sin y + y \cos y)]$$

$$\begin{aligned}
 f'(z) &= e^x [\cos y + i \sin y + x (\cos y + i \sin y) - y (\sin y - i \cos y)] \\
 &= e^x [e^{iy} + x e^{iy} - y (\sin y - i \cos y)]
 \end{aligned}$$

$$\begin{aligned}
 \text{ii)} \quad f(z) &= \sinh z = \frac{e^z - e^{-z}}{2} \\
 &= \frac{1}{2} \{ e^{x+iy} - e^{-(x+iy)} \} \\
 &= \frac{1}{2} \{ e^x \cdot e^{iy} - e^{-x} \cdot e^{-iy} \} \\
 &= \frac{1}{2} \{ e^x [\cos y + i \sin y] - e^{-x} [\cos y - i \sin y] \} \\
 &= \frac{1}{2} \{ (\cos y + i \sin y) (e^x - e^{-x}) \} \\
 &= \frac{(e^x - e^{-x}) \cos y}{2} + i \frac{(e^x - e^{-x}) \sin y}{2}
 \end{aligned}$$

$$\begin{aligned}
 (X) \quad f(z) &= \sinh x \cos y + i \sinh x \sin y \\
 \Rightarrow u &= \sinh x \cos y \\
 v &= \sinh x \sin y \\
 \frac{\partial u}{\partial x} &= \cos y \cosh x & \frac{\partial u}{\partial y} &= -\sinh x \sin y \\
 \frac{\partial v}{\partial x} &= \cosh x \sin y & \frac{\partial v}{\partial y} &= \sinh x \cos y
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \{ e^x [\cos y + i \sin y] - e^{-x} [\cos y - i \sin y] \} \\
 &= \frac{1}{2} \{ (e^x - e^{-x}) \cos y + i (e^x + e^{-x}) \sin y \} \\
 &= \sinh x \cos y + i \cosh x \sin y \\
 &= u + iv
 \end{aligned}$$

$$\begin{aligned}
 u &= \sinh x \cos y & v &= \cosh x \sin y \\
 \frac{\partial u}{\partial x} &= \cosh x \cos y & \frac{\partial v}{\partial x} &= \sinh x \sin y \\
 \frac{\partial u}{\partial y} &= -\sinh x \sin y & \frac{\partial v}{\partial y} &= \cosh x \cos y
 \end{aligned}$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

CR eqn^s hold good

$\therefore f(z) = \sinh z$ is Analytic functⁿ.

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$= \cosh x \cos y + i \sinh x \sin y$$

$$= \frac{(e^x + e^{-x})}{2} \cos\left(\frac{e^{iy} + e^{-iy}}{2}\right) + i \left(\frac{e^x - e^{-x}}{2}\right) \left(\frac{e^{iy} - e^{-iy}}{2}\right)$$

put $x=z$ & $y=0$

$$\boxed{f'(z) = \cosh x}$$

iv) $f(z) = \log z \{ = \log(x+iy) \text{ can't use } \}$ $\therefore$ convert to polar.

In polar form $z = re^{i\theta}$

$$f(z) = \log(re^{i\theta}) = \log r + \log e^{i\theta} = \log r + i\theta$$

$$= u + i v$$

$$u = \log r \quad \theta = v$$

$$\frac{\partial u}{\partial r} = \frac{1}{r} \quad \frac{\partial u}{\partial \theta} = 0$$

$$\frac{\partial v}{\partial r} = 0 \quad \frac{\partial v}{\partial \theta} = 1$$

$$\text{since } \frac{1}{r} \frac{\partial v}{\partial \theta} = \frac{1}{r} \frac{\partial u}{\partial \theta} \quad \& \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$$

CR eq^{ns} are satisfied $\therefore \log z$ is analytic.

$$f'(z) = e^{-i\theta} \left[\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right]$$

$$= e^{-i\theta} \left[\frac{1}{r} + i \cdot 0 \right] = \frac{e^{-i\theta}}{r} = \frac{1}{z}$$

$$v) f(z) = z^n$$

using polar form $\rightarrow z = re^{i\theta}$

$$f(z) = (re^{i\theta})^n = r^n e^{in\theta}$$

$$= r^n [\cos n\theta + i \sin n\theta]$$

$$u = r^n \cos n\theta \text{ and } v = r^n \sin n\theta$$

$$\frac{\partial u}{\partial r} = n r^{n-1} \cos n\theta \quad \frac{\partial v}{\partial r} = n r^{n-1} \sin n\theta$$

$$\frac{\partial u}{\partial \theta} = -r^n \sin n\theta \quad \frac{\partial v}{\partial \theta} = r^n \cos n\theta$$

$$\frac{\partial u}{\partial r} = n r^{n-1} \cos n\theta = \frac{1}{r} \frac{\partial v}{\partial \theta}$$

$$\frac{\partial v}{\partial r} = n r^{n-1} \sin n\theta = -\frac{1}{r} \frac{\partial u}{\partial \theta}$$

$$f'(z) = e^{i\theta} \left[\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right]$$

$$= e^{i\theta} [n r^{n-1} \cos n\theta + i n r^{n-1} \sin n\theta]$$

$$= e^{i\theta} [n r^{n-1} (e^{in\theta})]$$

$$= n r^{n-1} e^{i(n\theta + \theta)}$$

$$= n r^{n-1} e^{i\theta(n+1)}$$

$$= n r^{n-1} e^{i\theta(n+1)}$$

$$= n (re^{i\theta})^{n-1}$$

$$= n z^{n-1}$$

Harmonic Functions-

A function ϕ is said to be harmonic if it satisfies laplace eqⁿ. $\nabla^2 \phi = 0$

In cartesian form $\phi(x, y)$ is harmonic if

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0$$

In polar form $\phi(r, \theta)$ is harmonic if

$$\frac{\partial^2 \phi}{\partial r^2} + \frac{1}{r} \frac{\partial \phi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \phi}{\partial \theta^2} = 0$$

REMARK: The real & imaginary parts of an analytic functⁿ are harmonic.

• ^{NOT} The converse is not true i.e. we can example for u & v satisfying laplace eqⁿ but not satisfying CR equations.

Analytic functⁿ $\xrightarrow{\quad}$ Harmonic functⁿ
 $\xleftarrow{\quad}$

Definition:

let $u(x,y)$ be harmonic function then a functⁿ $v(x,y)$ is said to be harmonic conjugate of $u(x,y)$ if it satisfies following conditions-

$\Rightarrow v(x,y)$ is harmonic

$\Rightarrow$ CR eq^{ns} holds good.

Statement of theorem:

"If $f(z) = u + iv$ is an analytic function in a domain D , then v is harmonic conjugate of u conversely if v is harmonic conjugate of u in domain D then $f(z) = u + iv$ is analytic functⁿ in domain D ."

Constructⁿ of analytic functⁿ $f(z)$ given it's real limg. parts by Milne-Thomson's Method:

Procedure:-

(I) Cartesian Form:-

Step 1: Given u & v as a function of x & y .

Step 2: Find partial derivatives $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$.

Step 3: Consider $f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$

Case (1) : If u is given then use CR equation-

$$\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

$$\therefore f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \quad \text{--- (1)}$$

Case ii) If V is given

then use CR equation $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$

$$f'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x} \quad \text{--- (1)}$$

Step 4: Substitute expressions for partial derivatives in RHS of (1) & (2) & then put $x=z$ & $y=0$ to obtain $f'(z)$ as a function of z only.

Step 5: Integrate wrt z to get $f(z)$.

(II) Polar Form:

Step 1: Given u & v as functⁿ of r & θ .

Step 2: Find partial derivatives $\frac{\partial u}{\partial r}, \frac{\partial v}{\partial r}, \frac{\partial u}{\partial \theta}, \frac{\partial v}{\partial \theta}$

Step 3: Consider $f'(z) = e^{-i\theta} \left[\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right]$

Case (i) if u is given: then using CR eqⁿ

$$\frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$$

$$\therefore f'(z) = \left(\frac{\partial u}{\partial r} - \frac{i}{r} \frac{\partial u}{\partial \theta} \right) e^{-i\theta} \quad \text{--- (1)}$$

Case (ii) if v is given

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}$$

$$\therefore f'(z) = e^{-i\theta} \left[\frac{1}{r} \frac{\partial v}{\partial \theta} + i \frac{\partial v}{\partial r} \right] \quad \text{--- (2)}$$

Step 4: Substitute expressions for partial derivatives in RHS of (1) & (2) & then put $r=z$ & $\theta=0$ to obtain $f'(z)$ as functⁿ of z only.

Step 5: Integrate wrt z to get $f(z)$.

QPH> Determine analytic functⁿ whose real part is.

i) $x \cos 2y - y \sin 2y$ {real part}

given $u = x \cos 2y - y \sin 2y$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

use CReqⁿ $\frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$

$$f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\frac{\partial u}{\partial x} = \cos 2y \quad \frac{\partial u}{\partial y} = -2x \sin 2y - 2y \cos 2y - \sin 2y$$

$$\therefore f'(z) = \cos 2y + i(2x \sin 2y + 2y \cos 2y + \sin 2y)$$

$$f'(z) = \cos 2y + \text{put } x = z, y = 0$$

$$f'(z) = \cos 2 \cdot 1 + i[2z(0) + 0] = 1$$

$$f'(z) = 1$$

Integrate wrt z

$$f(z) = z + c$$

ii) $\cos x \cosh y$

given $u = \cos x \cosh y$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\frac{\partial u}{\partial x} = -\sin x \cosh y \quad \frac{\partial u}{\partial y} = \cos x \sinh y$$

$$f'(z) = -\sin x \cosh y + i \cos x \sinh y$$

put $x = z$ & $y = 0$

$$f'(z) = -\sin z - i \cos z(0)$$

$$\{\cosh y = 1, \sinh y = 0\}$$

$$= -\sin z$$

Integrate wrt z

$$f(z) = \cos z + C //$$

iii) $e^x \cos y$

given $u = e^x \cos y$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\frac{\partial u}{\partial x} = e^x \cos y \quad \frac{\partial u}{\partial y} = -e^x \sin y$$

$$f'(z) = e^x \cos y - i e^x \sin y$$

put $x=z$ & $y=0$

$$f'(z) = e^z$$

Integrate:

$$f(z) = e^z + C$$

LP5) Determine analytic functⁿ whose img. part is-

i) $\log(x^2 + y^2) + (x - 2y)$

ii) $e^x \{ (x^2 - y^2) \cos y - 2xy \sin y \}$

iii) $e^x (x \sin y + y \cos y)$

i) Given $v = \log(x^2 + y^2) + (x - 2y)$

$$f'(z) = \frac{\partial v}{\partial x} + i \frac{\partial v}{\partial y}$$

$$\text{CR eqⁿ } \frac{\partial v}{\partial y} = + \frac{\partial u}{\partial x}$$

$$f'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x} \quad \text{--- (1)}$$

$$\frac{\partial v}{\partial x} = \frac{2x}{x^2 + y^2} + 1 \quad \frac{\partial v}{\partial y} = \frac{2y}{x^2 + y^2} - 2$$

$$f'(z) = \frac{2y}{x^2 + y^2} - 2 + i \left(\frac{2x}{x^2 + y^2} + 1 \right)$$

put $x=z$ & $y=0$

$$f'(z) = -2 + i \left[\frac{2z}{z^2} + 1 \right]$$

integrate wrt:

$$f(z) = -2z + i [2 \log z + z] + C$$

ii) $v = e^x \{ (x^2 - y^2) \cos y - 2xy \sin y \}$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$= \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\frac{\partial v}{\partial y} = e^x \{ [(-2y) \cos y + \sin y (x^2 - y^2)] - \partial \sin y (x) - 2xy \cos y \}$$

$$\frac{\partial v}{\partial x} = e^x \{ 2x \cos y - 2y \sin y \} + \{ (x^2 - y^2) \cos y - 2xy \sin y \}$$

$$\therefore f'(z) = e^x [-2y \cos y - \sin y (x^2 - y^2) - 2x \sin y - 2xy \cos y] + i e^x [2x \cos y - 2y \sin y - 2xy \sin y + (x^2 - y^2) \cos y]$$

put $x=z$ & $y=0$

$$f'(z) = e^z [0] + i e^x [2z + z^2]$$

$$f'(z) = i e^z (2z + z^2)$$

integrate wrt z

$$f(z) = \int [e^z (2z^2 + z^3)]$$

$$f'(z) = i [(2z + z^2) e^z] - (2 + 2z) e^z + (2) e^z + C$$

$$= i [2ze^z + z^2 e^z - 2e^z - 2ze^z + 2e^z] + C$$

$$= i [2e^z - 2ze^z + z^2 e^z] + C$$

$$= i (z^2 e^z) + C //$$

iii) $e^x (x \sin y + y \cos y) = v$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

iii) $e^x \cos y$

given $u = e^x \cos y$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y}$$

$$\frac{\partial u}{\partial x} = e^x \cos y \quad \frac{\partial u}{\partial y} = -e^x \sin y$$

$$f'(z) = e^x \cos y - i e^x \sin y$$

put $x = z$ & $y = 0$

$$f'(z) = e^z$$

Integrate:

$$f(z) = e^z + C$$

LP5) Determine analytic functⁿ whose img. part is-

i) $\log(x^2 + y^2) + (x - 2y)$

ii) $e^x \{ (x^2 - y^2) \cos y - 2xy \sin y \}$

iii) $e^x (x \sin y + y \cos y)$

i) Given $v = \log(x^2 + y^2) + (x - 2y)$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\text{CR eq}^n \quad \frac{\partial v}{\partial y} = + \frac{\partial u}{\partial x}$$

$$f'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x} \quad \text{--- (1)}$$

$$\frac{\partial v}{\partial x} = \frac{2x}{x^2 + y^2} + 1 \quad \frac{\partial v}{\partial y} = \frac{2y}{x^2 + y^2} - 2$$

$$f'(z) = \frac{2y}{x^2 + y^2} - 2 + i \left(\frac{2x}{x^2 + y^2} + 1 \right)$$

put $x = z$ & $y = 0$

$$f'(z) = -2 + i \left[\frac{2z}{z^2} + 1 \right]$$

Integrate wrt:

$$f(z) = -2z + i [2 \log_e z + z] + C$$

ii) $V = e^x \{ (x^2 - y^2) \cos y - 2xy \sin y \}$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$= \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

$$\frac{\partial v}{\partial y} = e^x \{ [(-2y) \cos y + \sin y (x^2 - y^2)] - 2 \sin y (x) - 2xy \cos y \}$$

$$\frac{\partial v}{\partial x} = e^x \{ 2x \cos y - 2y \sin y \} + \{ (x^2 - y^2) \cos y - 2xy \sin y \}$$

$$\therefore f'(z) = e^x [-2y \cos y - \sin y (x^2 - y^2) - 2x \sin y - 2xy \cos y] + i e^x [2x \cos y - 2y \sin y - 2xy \sin y + (x^2 - y^2) \cos y]$$

put $x=z$ & $y=0$

$$f'(z) = e^z [0] + i e^x [2z + z^2]$$

$$f'(z) = i e^x (2z + z^2)$$

integrate wrt z

$$f(z) = i \left[e^z (2z^2 + \frac{z^3}{3}) \right]$$

$$f'(z) = i [(2z + z^2) e^z] - (2 + 2z) e^z + (2) e^z + C$$

$$= i [2ze^z + z^2 e^z - 2e^z - 2ze^z + 2e^z] + C$$

$$= i [2\cancel{ze^z} - 2\cancel{ze^z} + z^2 e^z] + C$$

$$= i (z^2 e^z) + C //$$

iii) $e^x (x \sin y + y \cos y) = v$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

LP9) If $f(z)$ is a regular functⁿ of z prove that

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) |f(z)|^2 = 4 |f'(z)|^2$$

— let $f(z) = u + iv$ is a regular functⁿ
 then $|f(z)| = \sqrt{u^2 + v^2}$
 $|f(z)|^2 = u^2 + v^2 = \phi$ (say) — ①

diff ① partially wrt x :

$$\frac{\partial \phi}{\partial x} = 2u \frac{\partial u}{\partial x} + 2v \frac{\partial v}{\partial x} \text{ — ②}$$

diff ② wrt x

$$\frac{\partial^2 \phi}{\partial x^2} = 2u \frac{\partial^2 u}{\partial x^2} + 2 \left(\frac{\partial u}{\partial x} \right)^2 + 2v \frac{\partial^2 v}{\partial x^2} + 2 \left(\frac{\partial v}{\partial x} \right)^2 \text{ — ③}$$

III¹⁴ diff ① partially wrt y twice: —

$$\frac{\partial \phi}{\partial y} = 2u \frac{\partial u}{\partial y} + 2v \frac{\partial v}{\partial y} \text{ — ④}$$

$$\frac{\partial^2 \phi}{\partial y^2} = 2u \frac{\partial^2 u}{\partial y^2} + 2 \left(\frac{\partial u}{\partial y} \right)^2 + 2v \frac{\partial^2 v}{\partial y^2} + 2 \left(\frac{\partial v}{\partial y} \right)^2 \text{ — ⑤}$$

Adding ③ & ⑤

$$\begin{aligned} \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} &= 2u \left[\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right] + 2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] \\ &\quad + 2v \left[\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} \right] + 2 \left[\left(\frac{\partial v}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial y} \right)^2 \right] \end{aligned}$$

Since $f(z) = u + iv$ is a regular functⁿ of z , u & v satisfy CR eq^{ns} & Laplace eq^{ns}.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ \& } \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

$$\text{and } \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \text{ \& } \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0$$

putting in eqⁿ ⑥

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0 + 0 + 2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] + 2 \left[\left(\frac{\partial v}{\partial y} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right]$$

$$\text{LHS} = 4 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] \quad \text{--- (7)}$$

Now: $f(z) = u + iv$

$$\Rightarrow f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$|f'(z)|^2 = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2$$

$$4 |f'(z)|^2 = 4 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] \quad \text{--- (8)}$$

LHS = RHS.

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \phi = 4 |f'(z)|^2$$

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) |f(z)| = 4 |f'(z)|^2 // \text{proved}$$

LP10) If z is harmonic holomorphic or analytic functⁿ of z . prove that $\left[\frac{\partial}{\partial x} |f(z)| \right]^2 + \left[\frac{\partial}{\partial y} |f(z)| \right]^2 = |f'(z)|^2$

proof: let $f(z) = u + iv$ is analytical functⁿ.

$$|f(z)| = \sqrt{u^2 + v^2}$$

$$|f(z)|^2 = u^2 + v^2 = \phi^2 \text{ (say)} \quad \text{--- (1)}$$

diff (1) wrt x partially

$$2\phi \frac{\partial \phi}{\partial x} = 2u \frac{\partial u}{\partial x} + 2v \frac{\partial v}{\partial x} \quad \text{--- (2)}$$

diff (1) wrt y partially

$$2\phi \frac{\partial \phi}{\partial y} = 2u \frac{\partial u}{\partial y} + 2v \frac{\partial v}{\partial y} \quad \text{--- (3)}$$

squaring & adding (2) & (3)

$$\phi^2 \left[\left(\frac{\partial \phi}{\partial x} \right)^2 + \left(\frac{\partial \phi}{\partial y} \right)^2 \right] = u^2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right]$$

$$+ 2uv \frac{\partial u}{\partial y} \cdot \frac{\partial v}{\partial y} + v^2 \left[\left(\frac{\partial v}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial y} \right)^2 \right] + 2uv \frac{\partial u}{\partial x} \cdot \frac{\partial v}{\partial x} \quad \text{--- (4)}$$

since $f(z)$ is analytic CR eq^{ns} are satisfied.

$$\text{i.e. } \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ \& } \frac{\partial v}{\partial x} = -\frac{\partial u}{\partial y}$$

eqⁿ (H) reduces to

$$\begin{aligned} & \phi^2 \left[\left(\frac{\partial \phi}{\partial x} \right)^2 + \left(\frac{\partial \phi}{\partial y} \right)^2 \right] = \\ & = u^2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] + 2uv \frac{\partial u}{\partial x} \frac{\partial v}{\partial x} \\ & \quad + v^2 \left[\left(-\frac{\partial v}{\partial x} \right)^2 \right] + v^2 \left(\frac{\partial u}{\partial x} \right)^2 - 2uv \frac{\partial u}{\partial x} \frac{\partial v}{\partial x} \\ & = u^2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] \\ & = (u^2 + v^2) \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 (u^2 + v^2) = \phi^2 \\ & = (u^2 + v^2) \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] = \phi^2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] \end{aligned}$$

$$\Rightarrow \phi^2 \left[\left(\frac{\partial \phi}{\partial x} \right)^2 + \left(\frac{\partial \phi}{\partial y} \right)^2 \right] = \phi^2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right]$$

$$\text{LHS} = \left[\left(\frac{\partial \phi}{\partial x} \right)^2 + \left(\frac{\partial \phi}{\partial y} \right)^2 \right] = \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right]$$

$$\text{we have } f(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$|f'(z)|^2 = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \quad \text{--- (6)}$$

from (5) & (6)

$$\frac{\partial}{\partial x} |f(z)|^2 + \frac{\partial}{\partial y} |f(z)|^2 = |f'(z)|^2 //$$